JOURNAL OF POLITICAL DISCOURSE
ISSN Print: 2992-2763 – ISSN Online: 2992-4618 | Vol. 3, No. 3 / September, 2025
https://www.jopd.com.ng

EXPLORING THE IMPACT OF NURSE BURNOUT ON PATIENT SAFETY AND CARE QUALITY IN ACUTE HEALTH CARE SETTINGS

Nwaduhu Beruck Oruchkwu
berucknwaduhu2017@gmail.com

Abstract
Nursing is one of the most demanding professions in the healthcare system, and nurses are often at the frontline of patient care, facing emotional, physical, and psychological challenges daily. Over time, these pressures can result in **burnout**, characterized by emotional exhaustion, depersonalization, and a reduced sense of personal accomplishment. This phenomenon not only affects nurses' well-being but also has profound implications for the quality and safety of patient care. This article examines the causes, consequences, and management strategies for nurse burnout within healthcare institutions. Drawing from theoretical frameworks such as Maslach's Burnout Theory and the Job Demand Resources Model, the study explores how organizational factors, workload, and emotional labor contribute to burnout and affect patient outcomes. The article emphasizes evidence-based strategies, including supportive leadership, resilience training, and institutional reforms, to mitigate burnout and promote optimal nursing performance. The paper concludes by highlighting the essential need for systemic interventions and policy support to safeguard nurses' mental health, enhance patient care quality, and strengthen global health systems.
Keywords: Nurse burnout, Patient care quality, Healthcare settings, Burnout prevention, Nursing management

Introduction

The nursing profession remains a cornerstone of global healthcare delivery, serving as the backbone of patient care and hospital functionality. Nurses constitute the largest proportion of healthcare professionals and play a vital role in maintaining the quality, safety, and continuity of care across all health service settings. Their responsibilities extend far beyond clinical duties; they encompass patient advocacy, health education, emotional support, and coordination of multidisciplinary care. Despite this critical role, nursing is widely recognized as one of the most demanding and stressful professions, with practitioners facing immense physical, emotional, and psychological burdens on a daily basis. In acute healthcare settings such as emergency departments, intensive care units, and surgical wards these pressures are magnified by high patient acuity, unpredictable workloads, and constant exposure to human suffering. Over time, such cumulative stressors predispose nurses to burnout, a syndrome characterized by emotional exhaustion, depersonalization, and a diminished sense of personal accomplishment (Maslach & Leiter, 2016).

Burnout among nurses has become a growing concern within contemporary healthcare discourse. It not only threatens the well-being and retention of nurses but also poses a direct risk to patient safety and the overall quality of healthcare delivery. Freudenberger (1974) was among the first scholars to describe burnout as a state of chronic emotional and physical depletion caused by persistent work-related stress. Maslach and Jackson (1981) later refined this concept into a multidimensional construct encompassing three central components: emotional exhaustion, depersonalization, and reduced personal accomplishment. Emotional exhaustion refers to the feeling of being emotionally over extended and depleted of one's emotional resources; depersonalization involves developing cynical or detached attitudes toward patients; and reduced personal accomplishment reflects a decline in one's sense of professional efficacy and achievement. In nursing practice, these

symptoms manifest in decreased empathy, irritability, frequent absenteeism, and declining motivation all of which undermine professional performance and patient care quality.

The increasing prevalence of burnout among nurses reflects broader systemic and organizational challenges within healthcare systems. Factors such as heavy workloads, chronic understaffing, inadequate managerial support, and limited opportunities for professional growth have been consistently identified as major contributors to burnout (Aiken et al., 2018; Laschinger & Fida, 2015). Nurses in many acute care settings are frequently required to handle complex clinical situations with minimal support, while simultaneously managing administrative tasks, maintaining documentation, and adhering to institutional protocols. The resulting imbalance between job demands and available resources creates an environment conducive to stress and fatigue (Demerouti et al., 2001). When nurses are stretched beyond their capacity, the quality and safety of patient care inevitably suffer, leading to increased rates of medical errors, poor communication, and lower patient satisfaction (Hall et al., 2016).

Moreover, the psychological toll of nursing cannot be understated. Nurses are repeatedly exposed to pain, suffering, and death circumstances that require not only technical competence but also emotional resilience and empathy. Figley (2017) refers to this phenomenon as "compassion fatigue," a condition that often precedes burnout. Compassion fatigue arises from the cumulative emotional residue of caring for traumatized or critically ill patients, leading to emotional numbness, detachment, and a gradual loss of empathy. In acute care units, where nurses must make quick, life-altering decisions under pressure, emotional depletion can have devastating consequences. The inability to manage these emotional demands effectively can erode professional judgment, increase clinical mistakes, and compromise patient outcomes.

The global COVID-19 pandemic further illuminated the fragility of healthcare systems and the vulnerability of frontline nurses. During the pandemic, nurses worldwide encountered unprecedented workloads, extended shifts, limited protective equipment, and frequent exposure to life-threatening infections. The combination of moral distress, fear, and uncertainty led to an alarming escalation in burnout levels, depression, and post-traumatic stress symptoms among healthcare workers (Shanafelt et al., 2021). Many nurses reported feeling unsupported by their institutions and undervalued by the public, resulting in emotional withdrawal and heightened intentions to leave the profession. These trends have serious implications for healthcare sustainability, as the global nursing shortage continues to intensify. According to the World Health Organization (2022), the world faces an estimated shortfall of nearly six million nurses, particularly in low- and middle-income countries. The burnout epidemic thus not only threatens individual well-being but also the stability and functionality of healthcare systems globally.

In addition to organizational and emotional factors, cultural and systemic dynamics play a significant role in shaping the experience of burnout among nurses. In many developing countries, nurses work in environments characterized by inadequate infrastructure, scarce medical supplies, and low remuneration. These conditions exacerbate stress and reduce motivation, leading to high turnover rates and low job satisfaction. Conversely, in high-income countries, technological advancements and bureaucratic demands often impose new forms of strain, such as digital overload and excessive administrative responsibilities. Despite these contextual differences, the underlying issue remains consistent: when nurses are overburdened and under-supported, both care quality and patient safety decline. Burnout among nurses, therefore, is not merely an occupational hazard but a systemic indicator of dysfunction within healthcare organizations.

From a patient safety perspective, the implications of nurse burnout are profound. Burned-out nurses are more likely to make clinical errors, demonstrate lapses in infection control, and fail to adhere to evidence-based care protocols (Hall et al., 2016). Emotional

exhaustion reduces attentiveness, while depersonalization undermines empathy and communication key components of effective patient care. Research consistently demonstrates a correlation between nurse burnout and adverse patient outcomes, including medication errors, falls, and mortality (Aiken et al., 2018). Moreover, patients often perceive the emotional detachment of burned-out nurses as a lack of compassion or professionalism, leading to diminished trust and satisfaction. This erosion of the nurse-patient relationship not only compromises clinical care but also negatively affects patient experience and recovery.

Burnout also carries organizational consequences that extend beyond individual practitioners. High levels of burnout contribute to absenteeism, turnover, and early retirement, which in turn exacerbate staffing shortages and increase institutional costs. The cycle of burnout and attrition creates a self-perpetuating challenge: as more nurses leave, the remaining staff face higher workloads, greater stress, and an increased risk of burnout. Consequently, healthcare organizations face escalating recruitment and training expenses, alongside declining productivity and morale. Effective management of burnout, therefore, is not only a moral and professional responsibility but also an economic imperative for healthcare institutions seeking sustainability.

Given the magnitude of these challenges, addressing nurse burnout requires a multidimensional approach that encompasses individual, organizational, and policy-level interventions. Individual strategies such as mindfulness training, resilience-building workshops, and self-care education can help nurses develop coping mechanisms to manage stress (West et al., 2020). However, such interventions are insufficient in isolation. Systemic reforms are needed to create supportive workplace cultures, implement fair staffing policies, and foster transformational leadership that empowers nurses and values their contributions (Laschinger & Fida, 2015). Transformational leadership, characterized by empathy, communication, and shared vision, has been shown to enhance nurse engagement and reduce emotional exhaustion. Likewise, institutional policies promoting work-life balance, psychological safety, and open communication can create environments where nurses thrive rather than merely survive.

In recent years, global health organizations and scholars have called for a paradigm shift in how healthcare institutions conceptualize and address burnout. Rather than viewing it solely as an individual problem requiring personal coping strategies, there is a growing recognition that burnout reflects deeper structural issues within healthcare systems. The Maslach Burnout Theory and the Job Demand–Resources Model (Demerouti et al., 2001) both provide valuable frameworks for understanding this relationship. These models suggest that burnout arises when organizational demands such as excessive workload, time pressure, and emotional labor overwhelm the available resources, including social support, autonomy, and professional recognition. In this sense, preventing burnout involves not just reducing stressors but also enhancing the resources that enable nurses to perform their duties effectively and meaningfully.

This article therefore seeks to contribute to the ongoing scholarly and policy discourse on nurse burnout by providing a comprehensive analysis of its causes, consequences, and management strategies. It highlights the complex interplay between personal, organizational, and systemic factors that contribute to burnout in acute healthcare settings. By integrating theoretical insights with empirical evidence, the study aims to illuminate pathways toward sustainable solutions that protect nurses' mental health while safeguarding patient safety and care quality. Ultimately, addressing nurse burnout is not merely an act of compassion ,it is an essential investment in the resilience, quality, and sustainability of healthcare systems worldwide.

Literature Review

Burnout among nurses has been widely recognized as a major occupational and public health issue that has profound consequences for healthcare systems across the world. It is a multidimensional phenomenon that arises from chronic workplace stress, emotional exhaustion, depersonalization, and a diminished sense of personal accomplishment (Maslach & Leiter, 2016). Over the decades, a significant body of literature has explored the antecedents, correlates, and consequences of nurse burnout, emphasizing its detrimental impact on both employee well-being and patient outcomes. The literature consistently demonstrates that burnout is not an isolated event but rather the result of an interplay between individual vulnerabilities, organizational factors, and systemic constraints that shape the nursing work environment (Aiken et al., 2018).

Conceptualizing Nurse Burnout

The concept of burnout was first introduced by Freudenberger (1974) to describe a state of emotional depletion and reduced motivation among individuals working in helping professions. Later, Maslach and Jackson (1981) provided a more systematic and multidimensional definition, identifying three key components: emotional exhaustion, depersonalization, and reduced personal accomplishment. Emotional exhaustion refers to feelings of being emotionally over extended and fatigued due to excessive demands; depersonalization represents a detached, cynical, or impersonal response toward recipients of care; while reduced personal accomplishment involves a negative evaluation of one's work and competence. Within nursing, these dimensions manifest as fatigue, irritability, loss of empathy, and declining work efficiency.

Burnout is now understood not merely as an individual psychological problem but as a workplace syndrome that arises when professional expectations exceed available resources and coping capacities (Maslach & Leiter, 2016). The literature emphasizes that nurses experience burnout differently depending on their clinical context, level of experience, and type of patient interaction. For example, nurses in intensive care units often report higher burnout levels due to the life-threatening nature of cases, frequent exposure to mortality, and constant need for rapid decision-making. Similarly, emergency room nurses encounter unpredictable workloads and exposure to trauma, which heighten stress and emotional fatigue.

Workload and Staffing Issues

One of the most widely studied and empirically validated determinants of nurse burnout is workload. Aiken et al. (2018) demonstrated that nurse-to-patient ratios significantly influence burnout levels and patient outcomes. When nurses are responsible for too many patients simultaneously, their ability to provide individualized, compassionate, and timely care declines. Excessive workload not only causes physical fatigue but also contributes to mental exhaustion and cognitive overload. This situation increases the likelihood of errors in medication administration, infection control, and patient monitoring, thereby compromising patient safety.

Insufficient staffing is another core driver of burnout and dissatisfaction. In healthcare environments with chronic understaffing, nurses are compelled to work long hours, often with limited breaks and inadequate support. Over time, this leads to a sense of helplessness and disengagement, as the effort required to meet patient needs becomes unmanageable. The pressure to meet performance indicators while handling high patient volumes can also create moral distress when nurses feel unable to deliver the quality of care they believe patients deserve. This moral distress accumulates into emotional exhaustion, a major component of burnout (Maslach & Leiter, 2016).

Workload and staffing challenges also interact with other organizational stressors such as shift work, mandatory overtime, and inconsistent scheduling. Rotating shifts disrupt circadian rhythms and sleep patterns, leading to chronic fatigue, irritability, and reduced concentration. Studies have consistently shown that nurses who work irregular shifts or extended hours are at greater risk of burnout, workplace injuries, and psychological strain. In acute healthcare settings, where patient acuity is high and clinical urgency is constant, these stressors are magnified, creating conditions that make burnout almost inevitable without adequate institutional support.

Organizational Culture and Leadership

The organizational environment in which nurses work plays a pivotal role in either exacerbating or mitigating burnout. Research underscores that organizational culture, leadership style, and communication patterns profoundly affect nurses' job satisfaction, motivation, and resilience (Laschinger & Fida, 2015). A supportive work culture characterized by open communication, mutual respect, and shared decision-making can serve as a buffer against stress, while authoritarian or unsupportive management can intensify it.

Laschinger and Fida (2015) highlight the importance of authentic and transformational leadership in fostering positive work environments that promote nurse engagement. When nurses perceive their leaders as empathetic, transparent, and empowering, they are more likely to feel valued and committed to their work. Transformational leaders inspire and motivate employees by articulating a shared vision, providing mentorship, and recognizing individual contributions. Conversely, when leadership fails to provide clear direction, constructive feedback, or recognition, nurses often experience feelings of isolation, frustration, and burnout.

Organizational culture also determines how institutions respond to stress and failure. In punitive environments where mistakes are met with blame rather than learning, nurses may suppress errors, leading to emotional distress and fear. Such cultures discourage teamwork and open communication, further contributing to burnout. In contrast, a culture of safety and support where nurses can discuss challenges and receive constructive feedback promotes psychological well-being and reduces depersonalization. Leadership development programs that emphasize emotional intelligence, communication skills, and conflict resolution have been shown to enhance both nurse satisfaction and patient care outcomes.

Emotional and Psychological Factors

The emotional demands of nursing are among the most profound contributors to burnout. Nurses frequently encounter pain, suffering, and death, often forming close emotional connections with patients and their families. Over time, the cumulative exposure to trauma and loss can result in compassion fatigue a precursor to burnout (Figley, 2017). Compassion fatigue manifests as emotional numbness, apathy, and difficulty empathizing with patients, undermining the quality of care and satisfaction derived from professional duties.

Figley (2017) describes compassion fatigue as "the cost of caring," emphasizing that it occurs when the energy devoted to helping others depletes one's emotional reserves. In nursing, this fatigue is intensified by ethical dilemmas and moral conflicts, such as witnessing preventable suffering, being unable to provide adequate care due to resource constraints, or experiencing repeated patient deaths. These stressors can lead to a loss of professional meaning and fulfillment, culminating in burnout.

Psychologically, burnout also interacts with personal traits such as perfectionism, empathy, and self-efficacy. Nurses who have high expectations of themselves and their professional roles may internalize organizational failures as personal shortcomings, increasing vulnerability to emotional exhaustion. Moreover, the stigma surrounding mental

health within healthcare professions often prevents nurses from seeking psychological support. As a result, burnout progresses unchecked until it manifests in physical illness, absenteeism, or career withdrawal.

Impact of Burnout on Patient Care and Safety

The literature consistently shows a direct and measurable relationship between nurse burnout and patient safety. Hall et al. (2016) in their systematic review found that burnout negatively affects patient care outcomes across diverse clinical settings. Nurses experiencing emotional exhaustion tend to have reduced attentiveness, slower response times, and poorer decision-making abilities. Depersonalization leads to diminished empathy and patient engagement, while reduced personal accomplishment correlates with decreased motivation to maintain quality standards.

Patients cared for by burned-out nurses are more likely to experience medication errors, hospital-acquired infections, and delayed treatment. Additionally, emotional detachment and poor communication can lead to misunderstandings between nurses and patients, resulting in decreased satisfaction and trust. Burnout also impairs teamwork and interprofessional collaboration, which are essential for coordinated and safe patient care.

From a systemic perspective, the cumulative effects of burnout extend beyond individual patient interactions. High burnout levels contribute to staff turnover, absenteeism, and decreased morale, which collectively weaken healthcare delivery systems. The shortage of experienced nurses not only disrupts continuity of care but also places additional pressure on remaining staff, perpetuating a cycle of overwork and emotional exhaustion. This dynamic illustrates how burnout is both a symptom and a cause of declining healthcare quality.

Preventive Strategies and Management Approaches

Given its multi factorial nature, preventing and managing nurse burnout requires comprehensive and evidence-based strategies. Scholars emphasize that interventions must address both individual and organizational levels to be effective. Individual strategies such as mindfulness, stress reduction techniques, and resilience training have gained prominence in recent years. These approaches focus on enhancing nurses' capacity to regulate emotions, manage stress, and maintain psychological balance under pressure (West et al., 2020). Mindfulness practices, for example, teach nurses to remain present and nonjudgmental, reducing emotional reactivity to stressful situations.

However, while personal coping mechanisms are valuable, they are insufficient if systemic issues remain unaddressed. Organizational interventions are equally crucial. Increasing staffing levels, ensuring fair workload distribution, and promoting flexible scheduling are key steps in reducing stress and fatigue. Moreover, creating peer support systems and counseling services allows nurses to process emotional experiences constructively. Leadership training for nurse managers can further foster environments of empathy, collaboration, and recognition factors shown to reduce burnout and enhance job satisfaction (Laschinger & Fida, 2015).

Institutional policies should also prioritize work-life balance by encouraging adequate rest periods, limiting overtime, and providing opportunities for professional development. Recognition and reward systems that acknowledge nurses' contributions can improve morale and reinforce a sense of purpose. As Aiken et al. (2018) demonstrated, when nurses perceive their workplaces as supportive and fair, both burnout and turnover rates decrease significantly.

Systemic and Policy-Level Considerations

Beyond institutional strategies, systemic reforms are necessary to address the structural determinants of burnout. Governments and health authorities play an essential role in setting standards for safe staffing ratios, regulating working hours, and funding wellness programs. Policy frameworks should emphasize the need for mental health support within healthcare settings and allocate resources for continuous education on stress management. The World Health Organization (2022) advocates for global investments in nursing education, leadership development, and workforce sustainability to prevent burnout and ensure high-quality care delivery.

The recognition of burnout as an occupational hazard necessitates its inclusion in occupational health and safety policies. Proactive monitoring of burnout indicators through periodic surveys and psychological assessments can help institutions identify at-risk staff early and implement preventive measures. Integrating technology to streamline administrative tasks, optimize workflow, and reduce documentation burden can also alleviate cognitive overload, allowing nurses to devote more time to direct patient care.

Research Objectives

This article aims to

Examine the key factors contributing to burnout among nurses in healthcare settings.
Assess the relationship between nurse burnout and the quality of patient care.
Identify evidence-based strategies to prevent and manage burnout in nursing practice.
Propose policy and institutional reforms that promote nurses' well-being and sustainable healthcare delivery.

Research Questions

The study is guided by the following research questions:

What are the primary causes of burnout among nurses?
How does burnout influence the quality and safety of patient care?
What preventive and management strategies can effectively reduce nurse burnout?
How can healthcare organizations foster supportive environments that enhance nurses' resilience and job satisfaction?

Theoretical Framework

This study draws on two main theoretical perspectives: **Maslach's Burnout Theory** and the **Job Demand Resources (JD-R) Model**.

Maslach's Burnout Theory posits that burnout arises from chronic workplace stressors that have not been successfully managed. The three dimensions emotional exhaustion, depersonalization, and reduced personal accomplishment are central to understanding how sustained stress undermines performance and well-being (Maslach & Leiter, 2016).

The Job Demand–Resources Model (Demerouti et al., 2001) explains burnout as a function of imbalance between job demands (workload, time pressure, emotional strain) and available resources (support, autonomy, feedback). When demands outweigh resources, energy depletion and disengagement occur. In the nursing context, this imbalance is common due to high patient loads and insufficient institutional support.

Together, these theories provide a framework for analyzing how structural and psychological factors interact to influence burnout and, consequently, patient care quality.

Research Methodology

This article employs a **qualitative descriptive design** using secondary data derived from scholarly journals, institutional reports, and empirical studies published between 2015 and

2024. The purpose is to synthesize current evidence on nurse burnout and identify effective interventions.

Data Sources
Peer-reviewed articles were retrieved from databases such as PubMed, ScienceDirect, and Google Scholar using keywords like "nurse burnout," "patient care quality," and "burnout prevention in nursing."

Inclusion Criteria
Studies were selected based on relevance, publication in reputable journals, and focus on professional nursing populations.

Data Analysis
Content analysis was employed to categorize findings into thematic areas,causes of burnout, effects on patient care and mitigation strategies. Patterns and relationships among variables were analyzed to identify common trends across studies.

Ethical Considerations
Because this study is based on secondary data, no direct ethical risks were involved. However, all sources were cited appropriately in line with academic integrity and the APA 7th edition citation style.

Conclusions
Nurse burnout is a critical and persistent problem that undermines the efficiency and humanity of healthcare systems worldwide. The findings indicate that burnout stems from excessive workload, poor management support, inadequate resources, and emotional fatigue. It leads to decreased quality of patient care, increased medical errors, and diminished job satisfaction among nurses.

Addressing burnout requires a holistic and systemic approach that includes institutional reforms, supportive leadership, and personal coping mechanisms. Prioritizing nurses' mental health is not merely an occupational issue it is a public health imperative. Enhancing nurse well-being directly correlates with improved patient safety and overall healthcare outcomes.

Recommendations
Healthcare institutions should establish supportive structures, including counseling services, mentorship programs, and fair workload distribution. Creating a culture of open communication and psychological safety can help nurses express concerns and seek help without fear of stigma.

Nurse managers and administrators must receive training in transformational leadership to promote empathy, collaboration, and empowerment. Leadership programs should focus on conflict resolution, team motivation, and stress reduction strategies.

Governments and professional bodies should enact policies mandating safe staffing ratios and regulating nurses' working hours. Policy measures should also include incentives for continued professional development and programs to reduce workplace violence.

Continuous education and resilience training should be incorporated into nursing curricula and workplace programs. Workshops on coping strategies, mindfulness, and emotional regulation can strengthen nurses' ability to manage stress effectively.

Regular mindfulness sessions, stress management workshops, and peer support groups should be institutionalized. Healthcare facilities should promote work-life balance through flexible scheduling, adequate rest periods, and recognition of nurses' contributions.

Motivation and retention can be improved through recognition of nurses' contributions, career advancement opportunities, and financial incentives. Providing access to professional counseling, mental health support, and wellness programs will further help in sustaining nurses' emotional and psychological well-being.

Institutional strategies should also focus on integrating technology to reduce administrative burdens, improve patient documentation, and optimize workflow efficiency. Collaboration between interdisciplinary teams can enhance patient care and reduce role-related stress for nurses.

Finally, periodic evaluation of nurse satisfaction and burnout indicators should be conducted to inform targeted interventions, ensuring continuous improvement of workplace conditions and patient care quality.

Major Contribution to Knowledge

This article contributes to the existing body of knowledge by integrating multiple perspectives on nurse burnout and its impact on patient care quality. It bridges theoretical understanding with practical strategies, offering a comprehensive framework for policymakers and healthcare administrators. The synthesis emphasizes that burnout is not solely an individual issue but an organizational and systemic one that demands proactive institutional responses.

The study also provides a foundation for future empirical research aimed at developing culturally sensitive and context-specific interventions for burnout prevention in low- and middle-income countries where the problem is often under-researched.

References

Aiken, L. H., Sloane, D. M., Ball, J., Bruyneel, L., Rafferty, A. M., & Griffiths, P. (2018). Patient satisfaction with hospital care and nurses in England: An observational study. *BMJ Open, 8*(1), e019189. https://doi.org/10.1136/bmjopen-2017-019189

Demerouti, E., Bakker, A. B., Nachreiner, F., & Schaufeli, W. B. (2001). The job demands–resources model of burnout. *Journal of Applied Psychology, 86*(3), 499–512.

Figley, C. R. (2017). Compassion fatigue: Coping with secondary traumatic stress disorder in those who treat the traumatized. *Routledge.*

Hall, L. H., Johnson, J., Watt, I., Tsipa, A., & O'Connor, D. B. (2016). Healthcare staff wellbeing, burnout, and patient safety: A systematic review. *PLOS ONE, 11*(7), e0159015.

Laschinger, H. K. S., & Fida, R. (2015). Linking nurses' perceptions of patient care quality to job satisfaction: The role of authentic leadership and burnout. *Journal of Nursing Management, 23*(6), 779–789.

Maslach, C., & Leiter, M. P. (2016). Understanding the burnout experience: Recent research and its implications for psychiatry. *World Psychiatry, 15*(2), 103–111.

Shanafelt, T., Ripp, J., & Trockel, M. (2021). Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. *JAMA, 323*(21), 2133–2134.

West, C. P., Dyrbye, L. N., & Shanafelt, T. D. (2020). Physician burnout: Contributors, consequences, and solutions. *Journal of Internal Medicine, 283*(6), 516–529.

World Health Organization (WHO). (2022). *State of the world's nursing 2022: Investing in education, jobs, and leadership.* Geneva: WHO.